

Expression analysis of genes encoding *TEX11*, *TEX12*, *TEX14* and *TEX15* in testis tissues of men with non-obstructive azoospermia

Parnaz Borjian Boroujeni¹, Marjan Sabbaghian², Mehdi Totonchi^{1,3}, Niloofar Sodeifi², Homa Sarkardeh¹, Azam Samadian³, Mohammad Ali Sadighi-Gilani^{2,4}, Hamid Gourabi¹

¹Department of Genetics, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

²Department of Andrology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran

³Department of Stem Cells and Developmental Biology, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran

⁴Department of Urology, Shariati Hospital, Tehran University of Medical Science, Tehran, Iran

Congresses where the study was presented:

Middle East Fertility Society (MEFS) congress 2012, 2013

Royan International Twin Congress 2013, 2014

ABSTRACT

Objective: Spermatogenesis is a complex process controlled by a plethora of genes. Changes in expression and function of these genes may thus lead to spermatogenic deficiency and male infertility. *TEX11*, *TEX12*, *TEX14* and *TEX15* are germ cell-specific genes expressed in the testis. *TEX11*, involved in the initiation and maintenance of chromosome synapses in meiotic chromosomes, has been shown to be essential for meiosis and fertility in males. *TEX14*, a component of intercellular bridges in germ cells, is required for spermatogenesis and fertility. *TEX12* and *TEX15* are essential for correct assembly of the synaptonemal complex and thus meiosis progression.

Methods: In order to examine whether changes in expression of these genes is associated with impaired spermatogenesis, expression levels of these genes were quantified by RT-qPCR on samples retrieved from infertile patients submitted to diagnostic testicular biopsy at Royan institute. Samples were divided into two groups of 18 patients with non-obstructive azoospermia considered as case; nine patients with obstructive azoospermia were included in the control group.

Results: A significant down-regulation of these genes was observed in the SCOS group when compared to the control group.

Conclusion: This result suggests that regular expression of *TEX11*, *TEX12*, *TEX14* and *TEX15* is essential for the early stages of spermatogenesis.

Keywords: Male infertility, Gene expression, Non-Obstructive Azoospermia, Intercellular bridges, Synaptonemal complex

INTRODUCTION

Male-factor infertility apparently accounts for 40% to 50% of infertile complications and may be defined by environmental reasons, infections, immunological or hormonal insufficiencies, while many are regulated by genetic factors (Hirsh, 2003; Brugh & Lipshultz, 2004; Boyle et al., 1992; Ma et al., 2016). Spermatogenesis is a complex process controlled by thousands of genes, where any change in the expression or function of these genes may lead to spermatogenic failure and male infertility (Zhou et al., 2009; Westerveld, 2008). Identification of stage-specific genes controlling spermatogenesis is thus important.

Previous studies have revealed that *TEX11*, *TEX12*, *TEX14* and *TEX15* expression is restricted to germ cells and is not detectable in somatic tissues of humans and mice (Wang et al., 2001; Adelman & Petrini, 2008). Multiple studies have shown that X-linked germ cell-specific genes such as testis-expressed gene 11 (*Tex11*) have significant roles in regulating male fertility (Zheng et al., 2010; Matzuk & Lamb, 2008). Furthermore, *TEX11* expression has been associated with the onset of spermatogenesis, although restricted to spermatocytes and round spermatids (Wang et al., 2005; Tang et al., 2011).

The synaptonemal complex (SC) is a large protein structure needed for synapsis and the successful completion of meiotic cell division. Incorrect assembly of this complex in mice results in maturation arrest and infertility (Bolcun-Filas et al., 2009; Hamer et al., 2008; Kouznetsova et al., 2011), and may similarly lead to infertility, recurrent miscarriage and aneuploidies such as Down syndrome in humans (de Vries et al., 2005; Gerton & Hawley, 2005; Page & Hawley, 2003). *Tex11*, *Tex12* and *Tex15* are needed in chromosomal synapsis and meiotic recombination. More specifically, *Tex11* forms distinct foci on homologous chromosomes that synapse with each other and seem to be a new component of meiotic nodules needed for recombination. In the absence of this protein, chromosomal synapsis stops and crossover formation is reduced (Stouffs & Lissens, 2012; Yang et al., 2008b). *Tex12* is a meiosis-specific protein essential for the progression of synapsis between homologous chromosomes in male and female germ cells (Hamer et al., 2008). Loss of *Tex15* leads to meiotic recombination failure, since this protein is responsible for the transfer of DNA repair proteins onto double strand break (DSB) locations (Yang et al., 2008a). *TEX12* is small with no known domains, but has orthologs in other mammals such as mice, cows, and dogs (Hamer et al., 2006; Wang et al., 2001). *TEX15* has orthologs in mammals and zebra fish (Yang et al., 2008a). It is abundantly expressed in post-meiotic germ cells, spermatogonia, and early spermatocytes, showing that this gene plays a role in different stages of spermatogenesis (Yang et al., 2008a; Wang et al., 2005). Interestingly, it is expressed in both the testis and ovaries, as is its mouse ortholog (Wang et al., 2001).

Germ cell intercellular bridges are required for fertility in invertebrates (Brill et al., 2000; Robinson et al., 1994; Robinson & Cooley, 1996) and have been preserved from invertebrates to humans (Fawcett et al., 1959), indicating their importance in fertility (Greenbaum et al., 2006;

2009). Mammalian intercellular bridges connect hundreds of germ cells in syncytia originated from a single spermatogonial stem cell (Huckins, 1971; Greenbaum *et al.*, 2006; 2009). In the absence of cell bridges in mammals, syncytium is thus not formed and as a result spermatogenesis is impaired (Greenbaum *et al.*, 2006; 2009). *TEX14* is needed to generate the germ cell intercellular bridges from the midbody (Greenbaum *et al.*, 2006). Human *TEX14* is preferentially expressed in the testis with its highest levels observed in spermatocytes and early round spermatids (Wu *et al.*, 2003). Greenbaum *et al.* (2006) showed that *Tex14*-/- mutant male mice do not form intercellular bridges, thus confirming its essentiality in mammalian spermatogenesis. It was later revealed that *Tex14* is also essential for complete spermatogenesis in Finnish Yorkshire boars as in mice (Sironen *et al.*, 2011). Chung *et al.* (2013) recognized new loci showing association with development of testicular germ cell cancer (TGCT), one of which (rs9905704) located in *TEX14*. It has also been shown that common genetic variations in *TEX14*, which is over-expressed in breast tumors, is associated with risk of breast cancer in Caucasians (Kelemen *et al.*, 2009). Recent studies showed that although *Tex11*-/-, *Tex14*-/- and *Tex15* -/- mutant male mice were sterile because of meiotic arrest and disturbance of spermatogenesis, mutant females were fertile. They suggested that the infertility observed in mutant male mice was caused by loss of function of these genes and that *Tex11*, *Tex14* and *Tex15* were needed for meiosis progression and fertility only in male mice (Yang *et al.*, 2008a; Greenbaum *et al.*, 2006; Sironen *et al.*, 2011). However, *TEX12* has identical localization patterns in oocytes and spermatocytes, revealing that this protein is a common part of the central element of the synaptonemal complex in mammals and is not sex-specific (Hamer *et al.*, 2006). *Tex12* -/- mutant male mice were azoospermic and therefore sterile because of failure in chromosomal synapsis progression, while *Tex12* -/- female mice were sterile because of loss of ovarian follicles (Hamer *et al.*, 2008).

The homology of amino acid sequences and expression patterns of the human and murine *TEX11*, *TEX12*, *TEX14* and *TEX15* genes (Wang *et al.*, 2001; Zheng *et al.*, 2010; Tang *et al.*, 2011; Stouffs *et al.*, 2009; Greenbaum *et al.*, 2006; Wu *et al.*, 2003; Hamer *et al.*, 2006 ; Yang *et al.*, 2008b) suggests that these genes may play some a few in human spermatogenesis. Given that deficiency of these genes results in azoospermic mice (Zheng *et al.*, 2010; Greenbaum *et al.*, 2006; Wu *et al.*, 2003; Hamer *et al.*, 2006; Yang *et al.*, 2008a), severe spermatogenic failure may thus be caused by their disruption. Therefore, this study aimed to replicate this finding in humans by examining the expression levels of *TEX11*, *TEX12*, *TEX14*, and *TEX15* in the testis tissue of patients with non-obstructive azoospermia and comparing them against controls.

MATERIAL AND METHODS

Patients

Twenty-seven tissue samples were obtained from 18 patients with non-obstructive azoospermia (NOA) and nine subjects with obstructive azoospermia. All participating infertile men had undergone testicular sperm extraction (TESE) procedures at the Royan Institute. All patients gave written informed consent and the Ethical Review Board of the Royan Institute approved the study (reference number EC/90/ 1050).

Nine of the 18 patients with NOA were diagnosed with Sertoli cell-only syndromes (SCOS), while the other nine were diagnosed with maturation arrest at the spermatocyte stage. The control group consisted of nine patients with obstructive azoospermia. Patients with abnormal

karyotypes and Y chromosome microdeletions were excluded from the study.

The age range of the patients at the time of diagnosis was 30 to 50 years and their mean age was 36.91 ± 5.39 years. Hormonal levels were measured using a competitive ELISA kit (Monobind, CA, USA) according to the procedure described by Zangeneh *et al.* (2015). Cytogenetic analysis was performed based on standard methods (Asia *et al.*, 2014).

Histological evaluation

After TESE, a portion of the testicular samples was submerged in Bouin's solution and sent for standard histopathological analysis (McLachlan *et al.*, 2007). Residual tissues were collected by the Royan Tissue Bank. Testicular histopathology was categorized according to the most recognized pattern of spermatogenesis process: samples with complete spermatogenesis (obstructive azoospermia), samples with spermatogenic maturation arrest at the spermatocyte stage or SCOS.

RNA extraction

Total RNA from frozen testis tissue was extracted with TRIzol (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. Approximately 50-100 mg of testis tissue was pipetted into one mL of TRIzol. RNA was detached with chloroform, precipitated with isopropanol, washed with 75% ethanol, and finally dissolved in DEPC treated water (Li *et al.*, 2006).

The concentration of RNA samples was determined spectrophotometrically by measuring absorption at 260 nm (Li *et al.*, 2006), while DNA and protein contamination were checked by optical density (OD) measurements at 260/280 nm.

The integrity of total RNA was evaluated by measuring 260:280 nm absorption ratios and by gel electrophoresis on 1.2% agarose gel (Yu *et al.*, 2007; Tang *et al.*, 2006).

DNase treatment and cDNA synthesis

Before total RNA reverse transcription (RT), the samples were treated with DNase to eliminate DNA contamination using the DNase I (RNase free) kit (Fermentas, Life Sciences, UK). Total RNA (2 μ g) was reverse-transcribed into cDNA in a reaction primed with random hexamer primers using the RevertAid H Minus First Strand cDNA Synthesis Kit (Fermentas, Life Sciences, UK) according to the manufacturer's recommendations. Polymerase chain reaction (PCR) primers for *TEX11*, *TEX12*, *TEX14*, and *TEX15* were designed using PerlPrimer v1.1.20 (Table 1).

After RNA extraction from testis tissue samples and cDNA synthesis, RT-PCR was performed with the related primers to confirm their expression in the samples of the three groups.

Quantitative RT-PCR (RT-qPCR)

RT-qPCR was carried out to confirm and analyze the expression levels of target genes in the SCOS, MA, and control groups. The reactions were processed on a 7500 Real Time PCR machine (Applied Biosystems, Carlsbad, CA, USA) using the Power SYBR Green PCR master mix (Applied Biosystem, EU). The amplification solution contained 10 μ l of Power SYBR Green PCR master mix, 50 ng of cDNA and 5 picomoles of each primer, yielding a final volume of 20 μ l. Cycling conditions included an initial step of enzyme activation at 95°C for 10 minutes, followed by 40 cycles of denaturation at 95°C for 15 seconds, annealing at 62°C for 30 seconds, and extension at 72°C for 30 seconds. Transcript levels normalized to human glyceraldehyde 3-phosphate dehydrogenase (GAPDH) showed minimum variation among individual samples. GAPDH was used as an internal

Table 1. Oligonucleotide primer sequences

| Gene | Forward primer | Reverse primer | Amplification temperature (°C) | Product size with cDNA (bp) | NCBI Reference sequence |
|-------|-------------------------|--------------------------|--------------------------------|-----------------------------|-------------------------|
| GAPDH | CTCATTCCTGGTATGAC AACGA | CTTCCTCTGTGCTTTGCT | 60 | 121 | NG-007073.2 |
| TEX11 | GCCTGAATAGAGCCCTTGTA | TAGATCAACTGCAACTGCCAT | 62 | 250 | NG-012574.1 |
| TEX12 | AGTCTCCAGTGCCAGATAGT | AGATTAATTCCCTTGCTCACATCA | 62 | 135 | NG-012574.1 |
| TEX14 | GGTTTATCCACCGCTCCCTC | CCTCTGCTCGCTTCAA | 62 | 198 | NG-012574.1 |
| TEX15 | AGGCAACATTCAAGCATCCA | AGTGAGCCAGGTAGTGATCTT | 62 | 141 | NG-012574.1 |

positive control. Each sample was run in duplicate and the mean value was calculated. No primer-dimer formation was observed during PCR amplification.

Immunohistochemistry

The expression of *TEX14* was analyzed in the SO/H ($n=3$), SCOS ($n=3$), and MA ($n=3$) groups by immunohistochemistry. First, testis tissue sections were deparaffinized in xylene washes and then rehydrated in descending concentrations of ethanol washes. Endogenous peroxidase activity was blocked for 1 h at 37°C with goat serum (Sigma-Aldrich, St Louis, MO, USA), and triton X 100 was used to permeabilize cell membranes. Primary (*Tex14* Antibody, NBP1-85424 Novus, Biologicals USA, 1:100) and secondary antibodies (goat anti-rabbit HRP, ab97051, 1:100) were added and protein localization was visualized using a Dako Liquid DAB+ Substrate Chromogen System (Dako, Glostrup, Denmark). Finally, cells were imaged using an Olympus IX 71 microscope (Julaton & Pera, 2011). Negative controls were generated by the same method as the positive controls; however, the primary antibody was replaced with PBS solution.

Statistical analysis

Data on clinical characteristics were shown as mean \pm SEM. The normality of variables was analyzed with the Kolmogorov test. Differences in the mean values among the three groups were analyzed by one-way analysis of variance (ANOVA). The tests cited above were performed on SPSS statistical software package (SPSS Inc, Chicago, IL, USA) version 22.0. Real-time data were processed and analyzed using a two-tailed t-test. Differences with p -values <0.05 were considered significant.

RESULTS

Patient clinical characteristics

The characteristics of patients including age and LH, FSH, and testosterone levels are listed in Table 2. There was no significant difference in age, LH or testosterone serum levels between the three groups; however, the FSH serum levels between these groups were significantly different ($p=0.01$).

Gene expression analyses

The RT-qPCR results demonstrated that transcripts of all four *TEX* genes existed in all samples of the SCOS, MA, and control groups. Differential expression analysis showed that expression of *TEX* genes is significantly lower in SCOS testes samples when compared with controls ($p<0.01$) (Figures 1-4). The expression of *TEX11* and *TEX14* was also significantly lower in MA samples compared with controls ($p\leq 0.038$) (Figures 1,3); however, expression of *TEX12* and *TEX15* was not significantly different (*TEX12,15 p>0.05*) (Figures 2,4). Expression levels of

TEX12, *TEX14* and *TEX15* in SCOS and MA samples were not significantly different (*TEX12, 14,15 p>0.05*) (Figures 2,3, and 4); however, *TEX11* expression was significantly lower in SCOS samples ($p=0.003$) (Figure 1).

Immunohistochemical analysis of *TEX14* expression

Immunohistochemical analysis with *TEX14* antibody indicated absence of Leydig cell staining in several testis samples in all three groups. *TEX14* was however detected in a small number of Sertoli cells, averaging 1-2 positive cells per tubule, consistent with the low level of RNA detected by RT-qPCR. The intensity of staining in germ cells of control tissues was higher than in MA tissues (Figure 5). *TEX14* was expressed specifically by germ cells at varying stages of spermatogenesis. Type A spermatogonia (SA), primary spermatocytes (S1), and early round spermatids (S3) were positively stained for *TEX14*. No positive signal was detected in SCOS tissue samples.

DISCUSSION

The same expression pattern of *TEX11*, *TEX12*, *TEX14* and *TEX15* genes in mice and humans indicates that these conserved genes may have an important role in the early stages of mammalian spermatogenesis (Zheng et al., 2010; Tang et al., 2011; Stouffs et al., 2009; Greenbaum et al., 2006; Wu et al., 2003; Hamer et al., 2006; Yang et al., 2008a). Although the function of these genes in humans is not well-known, their roles in the progression of mouse spermatogenesis (Zheng et al., 2010; Greenbaum et al., 2006; Wu et al., 2003; Hamer et al., 2006; Yang et al., 2008b) led us to hypothesize that they were likely to have the same roles in humans. To test this hypothesis, we compared the expression levels of these genes in testicular tissue samples of infertile men with two distinct histological patterns of spermatogenic failure (MA and SCOS) - our case group - to the levels observed in a control group including men with complete spermatogenesis (obstructive azoospermia).

Given that there was no germ cell in the testicular tissue samples of SCOS patients and all four *TEX* genes are germ cell-specific, down-regulation of these genes was expected. This observation is in line with the findings in mice, in which the lack of these genes disrupts the development of meiosis, resulting in male infertility (Zheng et al., 2010; Greenbaum et al., 2006; Wu et al., 2003; Hamer et al., 2006; Yang et al., 2008). The similar expression pattern and sequence conservation of *TEX11* seen in mice, humans, and pigs indicate that this gene is highly conserved and may have an important role in mammalian testicular function, including spermatogenesis (Zheng et al., 2010; Tang et al., 2011; Stouffs et al., 2009). *TEX11* expression in testis is correlated with the onset of spermatogenesis and is restricted to spermatocytes and round spermatids (Tang et al., 2011). Stage-specific expression of *TEX11* in

Table 2. Clinical characteristics of patient groups. Values are expressed as mean \pm SEM

| Patients groups | Age(years) | FSH(mIU/mL) | LH(mIU/mL) | Testosterone (ng/mL) |
|-----------------------------------|-------------------------|------------------|-----------------|----------------------|
| Maturation arrest (MA) | 37.28 \pm 3.28(30-50) | 5.62 \pm 1.51 | 4.32 \pm 1.26 | 5.28 \pm 0.84 |
| Sertoli-cell-only syndrome (SCOS) | 37.33 \pm 1.20(33-42) | 18.57 \pm 3.90 | 3.73 \pm 1.82 | 3.27 \pm 0.59 |
| Control | 36.33 \pm 1.24(30-42) | 8.37 \pm 1.73 | 3.70 \pm 0.87 | 5.61 \pm 3.57 |
| p-value | 0.924 | 0.010* | 0.941 | 0.237 |

Values are expressed as mean \pm SEM.

*Significant difference based on ANOVA.

Normal range of FSH: 1.5-12.4 mIU/mL

Normal range of LH: 1.0-10.0 mIU/mL

Normal range of Testosterone: 2.0-8.0ng/mL




Figure 1. Comparison of the expression levels of TEX11 between MA, SCOS, and control patients. *** $p<0.05$.




Figure 4. Comparison of the expression levels of TEX15 between MA, SCOS, and control patients. *** $p<0.05$.




Figure 2. Comparison of the expression levels of TEX12 between MA, SCOS, and control patients. *** $p<0.05$.




Figure 3. Comparison of the expression levels of TEX14 between MA, SCOS, and control patients. *** $p<0.05$.

porcine meiotic germ cells is in agreement with findings in mice, where the lack of *Tex11* disturbs the progression of meiosis and thus results in infertility. The loss of function of *Tex11* in *Tex11*-null mice resulted in meiotic arrest and deletion of spermatocytes (Yang *et al.*, 2008b). In summary, the expression pattern of *Tex11* is highly preserved in rodents and higher mammals such as pigs and humans (Tang *et al.*, 2011).

Yang *et al.* (2015) demonstrated that the frequency of rare *TEX11* mutations is significantly higher in azoospermic men, suggesting that *TEX11* is essential for human spermatogenesis and mutations in this single X-linked gene is the cause of infertility in \sim 1% of azoospermic men. The authors also reported that the level of *TEX11* protein must be above a critical threshold for meiosis to progress, and that low-expressing *TEX11* alleles may thus result in human male infertility. Yatsenko *et al.* (2015) hypothesized that mutations in human *TEX11* disrupt the formation and function of the synaptonemal complex, resulting in disturbance of pachytene synapsis, meiotic arrest, and azoospermia. The authors reported that hemizygous *TEX11* mutations were a common cause of meiotic arrest and azoospermia in infertile men.

Our results showed that *TEX11* expression was significantly decreased in MA patients when compared to controls. As dysfunction of *TEX11* results in meiotic arrest and sterility in mice (Adelman & Petrini, 2008; Yang *et al.*, 2008b; Stouffs *et al.*, 2009), impairment of spermatogenesis and maturation arrest in this group may be linked to *TEX11* down-regulation. This down-regulation was significantly more pronounced in SCOS patients than in patients with MA. Our results confirmed the involvement of *TEX11* in human spermatogenesis. Furthermore, our findings indicated that expression of this gene is required for the completion of spermatogenesis.




Figure 5. (A–D) Immunohistochemical analysis of adult testis sections with TEX14 antibody. (A) TEX14 is expressed in human testis cross-sections in obstructive azoospermia tissue samples. TEX14 is expressed specifically by germ cells at varying stages of spermatogenesis. Type A spermatogonia (SA), primary spermatocytes (S1) and early round spermatids (S3) stain positively for TEX14. (B) TEX14 is expressed in human testis cross-sections in MA tissue samples. (C) Negative control of testis cross-sections in MA tissue samples. (D) No positive signal is detected in adult human testis of SCOS tissue samples. The scale bar represents 10µm.

Tex12 expression is limited to meiotic cell division and its promoter, like many other germ cell-specific genes, is quenched in somatic cells by transcription factor E2F6 (Pohlers *et al.*, 2005). Previous studies showed that *TEX12* localizes to the central element of the synaptonemal complex. Their results also revealed that without *TEX12*, the synaptonemal complex lacks a correct central element structure, synapsis cannot complete, and meiotic recombination and crossing-over do not occur (Hamer *et al.*, 2006). Since loss of *TEX12* results in the inaccurate assembly of the synaptonemal complex and non-chromosomal

synapsis (Hamer *et al.*, 2006), reduced expression of this gene in the MA group in relation to the control group was expected; however, our results did not reveal a significant difference in expression between these two groups.

Previous studies of *Tex15* in mice showed this gene is necessary for chromosomal synapsis, DSB repair, and meiotic recombination during meiosis; therefore, *TEX15* is thought of as a possible factor in spermatogenic failure risk (Yang *et al.*, 2008a). Okutman *et al.* (2015) showed that a nonsense mutation in *TEX15* is the cause of spermatogenic defect in familial cases of teratozoospermia and

infertility. Immunohistochemistry confirmed these findings showing high levels of expression in germ cells and low levels of expression in Sertoli cells. The tests also revealed that *Tex15* knock-out produced spermatogenesis meiotic arrest and severe reduction in testicular size. Therefore, we expected to observe down-regulation of *TEX15* in the MA group when compared to controls. However, our results did not show any significant difference. This may possibly be due to high levels of expression in spermatogonia and early spermatocytes in the testicular tissue of MA patients. Previous studies have shown spermatogenic disruption in *Tex14* knockout mice before the first meiotic division due to loss of intercellular bridges (Bolcun-Filas *et al.*, 2009; Greenbaum *et al.*, 2006; 2009). Moreover, a study on *Tex14* mutant pigs showed that *Tex14* is also essential for the completion of spermatogenesis in pigs (Sironen *et al.*, 2011). Although *TEX14* localizes to intercellular bridges of both female and male mice germ cells, disruption of this gene leads to sterility only in males (Brill *et al.*, 2000; Greenbaum *et al.*, 2006). The comparison of *TEX14* expression between control and MA groups showed that *TEX14* expression had a significant decrease in MA samples. As dysfunction of *TEX14* results in meiotic arrest and sterility in mice (Greenbaum *et al.*, 2006; 2009; Wu *et al.*, 2003), impairment of spermatogenesis and maturation arrest may be linked to the down-regulation of *TEX14*. These results were confirmed by immunohistochemistry, with *TEX14* expressed in type A spermatogonia, primary spermatocytes, and early round spermatids in control samples. Conversely, this protein was absent or hardly visible in type A spermatogonia, primary spermatocytes, and early round spermatids of MA samples. These results support the possibility that *TEX14* under-expression in MA patients is etiologic, suggesting that the low expression of this protein in these patients is unlikely to be caused by the lack of germ cells.

CONCLUSION

This study demonstrated that *TEX11*, *TEX12*, *TEX14* and *TEX15* are essential for the completion of spermatogenesis. Down-regulation of these genes, especially *TEX14* and *TEX11*, may lead to impaired spermatogenesis in infertile men. Studies on other independent sample sets are required to confirm this association.

ACKNOWLEDGMENT

This study was funded by a grant from the Royan Institute, Tehran, Iran. The authors would like to thank to Dr. A.H. Shahverdi, Dr. A. Vosough, and Mrs. N. Fatemi for their invaluable support.

CONFLICTS OF INTEREST

None to declare

Corresponding Author:

Hamid Gourabi
Reproductive Biomedicine Research Center
Royan Institute for Reproductive Biomedicine
Tehran, Iran
E-mail: gourabi@royaninstitute.org

REFERENCES

- Adelman CA, Petrini JH. ZIP4H (*TEX11*) deficiency in the mouse impairs meiotic double strand break repair and the regulation of crossing over. *PLoS Genet.* 2008;4:e1000042. PMID: 18369460 DOI: 10.1371/journal.pgen.1000042
- Asia S, Vaziri Nasab H, Sabbaghian M, Kalantari H, Zari Moradi S, Gourabi H, Mohseni Meybodi A. A Rare De novo Complex Chromosomal Rearrangement (CCR) Involving Four Chromosomes in An Oligo-asthenosperm Infertile Man. *Cell J.* 2014;16:377-82. PMID: 24611143
- Bolcun-Filas E, Speed R, Taggart M, Grey C, de Massy B, Benavente R, Cooke HJ. Mutation of the mouse *Syce1* gene disrupts synapsis and suggests a link between synaptonemal complex structural components and DNA repair. *PLoS Genet.* 2009;5:e1000393. PMID: 19247432 DOI: 10.1371/journal.pgen.1000393
- Boyle CA, Khouri MJ, Katz DF, Annest JL, Kresnow MJ, DeStefano F, Schrader SM. The relation of computer-based measures of sperm morphology and motility to male infertility. *Epidemiology.* 1992;3:239-46. PMID: 1591323 DOI: 10.1097/00001648-199205000-00009
- Brill JA, Hime GR, Scharer-Schuksz M, Fuller MT. A phospholipid kinase regulates actin organization and intercellular bridge formation during germline cytokinesis. *Development.* 2000;127:3855-64. PMID: 10934029
- Brugh VM 3rd, Lipshultz LI. Male factor infertility: evaluation and management. *Med Clin North Am.* 2004;88:367-85. PMID: 15049583 DOI: 10.1016/S0025-7125(03)00150-0
- Chung CC, Kanetsky PA, Wang Z, Hildebrandt MA, Koster R, Skotheim RI, Kratz CP, Turnbull C, Cortessis VK, Bakken AC, Bishop DT, Cook MB, Erickson RL, Fosså SD, Jacobs KB, Korde LA, Kraggerud SM, Lothe RA, Loud JT, Rahman N, Skinner EC, Thomas DC, Wu X, Yeager M, Schumacher FR, Greene MH, Schwartz SM, McGlynn KA, Chanock SJ, Nathanson KL. Meta-analysis identifies four new loci associated with testicular germ cell tumor. *Nat Genet.* 2013;45:680-5. PMID: 23666239 DOI: 10.1038/ng.2634
- de Vries FAT, de Boer, E, van den Bosch, M, Baarens, WM, Ooms, M, Yuan L, Liu JG, van Zeeland AA, Heyting C, Pastink A. Mouse *Sycp1* functions in synaptonemal complex assembly, meiotic recombination, and XY body formation. *Genes Dev.* 2005;19:1376-89. PMID: 15937223 DOI: 10.1101/gad.329705
- Fawcett DW, Ito S, Slatterback D. The occurrence of intercellular bridges in groups of cells exhibiting synchronous differentiation. *J Biophys Biochem Cytol.* 1959;5:453-60. PMID: 13664686 DOI: 10.1083/jcb.5.3.453
- Gerton JL, Hawley RS. Homologous chromosome interactions in meiosis: diversity amidst conservation. *Nat Rev Genet.* 2005;6:477-87. PMID: 15931171 DOI: 10.1038/nrg1614
- Greenbaum MP, Iwamori N, Agno JE, Matzuk MM. Mouse *TEX14* is required for embryonic germ cell intercellular bridges but not female fertility. *Biol Reprod.* 2009;80:449-57. PMID: 19020301 DOI: 10.1095/biol-reprod.108.070649
- Greenbaum MP, Yan W, Wu MH, Lin YN, Agno JE, Sharma M, Braun RE, Rajkovic A, Matzuk MM. *TEX14* is essential for intercellular bridges and fertility in male mice. *Proc Natl Acad Sci U S A.* 2006;103:4982-7. PMID: 16549803 DOI: 10.1073/pnas.0505123103

- Hamer G, Gell K, Kouznetsova A, Novak I, Benavente R, Höög C. Characterization of a novel meiosis-specific protein within the central element of the synaptonemal complex. *J Cell Sci.* 2006;119:4025-32. PMID: 16968740 DOI: 10.1242/jcs.03182
- Hamer G, Wang H, Bolcun-Filas E, Cooke HJ, Benavente R, Höög C. Progression of meiotic recombination requires structural maturation of the central element of the synaptonemal complex. *J Cell Sci.* 2008;121:2445-51. PMID: 18611960 DOI: 10.1242/jcs.033233
- Hirsh A. Male subfertility. *BMJ.* 2003;327:669-72. PMID: 14500443 DOI: 10.1136/bmj.327.7416.669
- Huckins C. The spermatogonial stem cell population in adult rats. I. Their morphology, proliferation and maturation. *Anat Rec.* 1971;169:533-57. PMID: 5550532 DOI: 10.1002/ar.1091690306
- Julaton VT, Reijo Pera RA. NANOS3 function in human germ cell development. *Hum Mol Genet.* 2011;20:2238-50 PMID: 21421998 DOI:10.1093/hmg/ddr114
- Kelemen LE, Wang X, Fredericksen ZS, Pankratz VS, Pharoah PD, Ahmed S, Dunning AM, Easton DF, Vierkant RA, Cerhan JR, Goode EL, Olson JE, Couch FJ. Genetic variation in the chromosome 17q23 amplicon and breast cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2009;18:1864-8. PMID: 19454617 DOI: 10.1158/1055-9965.EPI-08-0486
- Kouznetsova A, Benavente R, Pastink A, Höög C. Meiosis in mice without a synaptonemal complex. *PLoS One.* 2011;6:e28255. PMID: 22164254 DOI: 10.1371/journal.pone.0028255
- Li HG, Liao AH, Ding XF, Zhou H, Xiong CL. The expression and significance of CATSPER1 in human testis and ejaculated spermatozoa. *Asian J Androl.* 2006;8:301-6. PMID: 16625279 DOI: 10.1111/j.1745-7262.2006.00132.x
- Ma TJ, Zhang XJ, Ding XP, Chen HH, Zhang YW, Ding M. Association of single nucleotide polymorphisms in UBR2 gene with idiopathic aspermia or oligospermia in Sichuan, China. *Andrologia.* 2016;48:1253-60. PMID: 26940145 DOI: 10.1111/and.12569
- Matzuk MM, Lamb DJ. The biology of infertility: research advances and clinical challenges. *Nat Med.* 2008;14:1197-213. PMID: 18989307 DOI: 10.1038/nm.f.1895
- McLachlan RI, Rajpert-De Meyts E, Hoei-Hansen CE, de Kretser DM, Skakkebaek NE. Histological evaluation of the human testis--approaches to optimizing the clinical value of the assessment: mini review. *Hum Reprod.* 2007;22:2-16. PMID: 16887924 DOI: 10.1093/humrep/del279
- Okutman O, Muller J, Baert Y, Serdarogullari M, Gultomruk M, Piton A, Rombaut C, Benkhaliha M, Teletin M, Skory V. Exome sequencing reveals a nonsense mutation in TEX15 causing spermatogenic failure in a Turkish family. *Hum Mol Genet.* 2015;24:5581-8. PMID: 26199321 DOI: 10.1093/hmg/ddv290
- Page SL, Hawley RS. Chromosome choreography: the meiotic ballet. *Science.* 2003;301:785-9. PMID: 12907787 DOI: 10.1126/science.1086605
- Pohlers M, Truss M, Frede U, Scholz A, Strehle M, Kuban RJ, Hoffmann B, Morkel M, Birchmeier C, Hagemeier C. A role for E2F6 in the restriction of male-germ-cell-specific gene expression. *Curr Biol.* 2005;15:1051-7. PMID: 15936277 DOI: 10.1016/j.cub.2005.04.060
- Robinson DN, Cant K, Cooley L. Morphogenesis of Drosophila ovarian ring canals. *Development.* 1994;120:2015-25. PMID: 7925006
- Robinson DN, Cooley L. Stable intercellular bridges in development: the cytoskeleton lining the tunnel. *Trends Cell Biol.* 1996;6:474-9. PMID: 15157506 DOI: 10.1016/0962-8924(96)84945-2
- Sironen A, Uimari P, Venhoranta H, Andersson M, Vilki J. An exonic insertion within Tex14 gene causes spermatogenic arrest in pigs. *BMC Genomics.* 2011;12:591. PMID: 22136159 DOI: 10.1186/1471-2164-12-591
- Stouffs K, Lissens W. X chromosomal mutations and spermatogenic failure. *Biochim Biophys Acta.* 2012;1822:1864-72. PMID: 22634129 DOI: 10.1016/j.bbadi.2012.05.012
- Stouffs K, Tournaye H, Liebaers I, Lissens W. Male infertility and the involvement of the X chromosome. *Hum Reprod Update.* 2009;15:623-37. PMID: 19515807 DOI: 10.1093/humupd/dmp023
- Tang A, Yu Z, Gui Y, Guo X, Long Y, Cai Z. Identification and characteristics of a novel testis-specific gene, Tsc24, in human and mice. *Biol Pharm Bull.* 2006;29:2187-91. PMID: 17077512 DOI: 10.1248/bpb.29.2187
- Tang L, Zeng W, Clark RK, Dobrinski I. Characterization of the porcine testis-expressed gene 11 (Tex11). *Spermatogenesis.* 2011;1:147-51. PMID: 22319663 DOI: 10.4161/spmg.1.2.16680
- Wang PJ, McCarrey JR, Yang F, Page DC. An abundance of X-linked genes expressed in spermatogonia. *Nat Genet.* 2001;27:422-6. PMID: 11279525 DOI: 10.1038/86927
- Wang PJ, Page DC, McCarrey JR. Differential expression of sex-linked and autosomal germ-cell-specific genes during spermatogenesis in the mouse. *Hum Mol Genet.* 2005;14:2911-8. PMID: 16118233 DOI: 10.1093/hmg/ddi322
- Westerveld GH. Unraveling the genetics of spermatogenic failure. [Thesis]. Amsterdam: University of Amsterdam; 2008. 128 p.
- Wu MH, Rajkovic A, Burns KH, Yan W, Lin YN, Matzuk MM. Sequence and expression of testis-expressed gene 14 (Tex14): a gene encoding a protein kinase preferentially expressed during spermatogenesis. *Gene Expr Patterns.* 2003;3:231-6. PMID: 12711554 DOI: 10.1016/S1567-133X(03)00036-X
- Yang F, Eckardt S, Leu NA, McLaughlin KJ, Wang PJ. Mouse TEX15 is essential for DNA double-strand break repair and chromosomal synapsis during male meiosis. *J Cell Biol.* 2008a;180:673-9. PMID: 18283110 DOI: 10.1083/jcb.200709057

Yang F, Gell K, van der Heijden GW, Eckardt S, Leu NA, Page DC, Benavente R, Her C, Höög C, McLaughlin KJ, Wang PJ. Meiotic failure in male mice lacking an X-linked factor. *Genes Dev.* 2008b;22:682-91. PMID: 18316482 DOI: 10.1101/gad.1613608

Yang F, Silber S, Leu NA, Oates RD, Marszalek JD, Skaltsky H, Brown LG, Rozen S, Page DC, Wang PJ. TEX11 is mutated in infertile men with azoospermia and regulates genome-wide recombination rates in mouse. *EMBO Mol Med.* 2015;7:1198-210. PMID: 26136358 DOI: 10.15252/emmm.201404967

Yatsenko AN, Georgiadis AP, Röpke A, Berman AJ, Jaffe T, Olszewska M, Westernströer B, Sanfilippo J, Kurpisz M, Rajkovic A, Yatsenko SA, Kliesch S, Schlatt S, Tüttelmann F. X-linked TEX11 mutations, meiotic arrest, and azoospermia in infertile men. *N Engl J Med.* 2015;372:2097-107. PMID: 25970010 DOI: 10.1056/NEJMoa1406192

Yu Z, Tang A, Gui Y, Guo X, Zhu H, Long Y, Li Z, Cai Z. Identification and characteristics of a novel testis-specific gene, Tsc21, in mice and human. *Mol Biol Rep.* 2007;34:127-34. PMID: 17091336 DOI: 10.1007/s11033-006-9026-6

Zangeneh F, Yazdi RS, Naghizadeh MM, Abedinia N. Effect of Ramadan Fasting on Stress Neurohormones in Women with Polycystic Ovary Syndrome. *J Family Reprod Health.* 2015;9:51-7. PMID: 26175759

Zheng K, Yang F, Wang PJ. Regulation of male fertility by X-linked genes. *J Androl.* 2010;31:79-85. PMID: 19875494 DOI: 10.2164/jandrol.109.008193

Zhou Y, Qin D, Tang A, Zhou D, Qin J, Yan B, Diao R, Jiang Z, Cai Z, Gui Y. Developmental expression pattern of a novel gene, TSG23/Tsg23, suggests a role in spermatogenesis. *Mol Hum Reprod.* 2009;15:223-30. PMID: 19240080 DOI: 10.1093/molehr/gap015